

## One-Sided Multivariate Tests for High Dimensional Data

Samruam Chongcharoen  
School of Applied Statistics,  
National Institute of Development Administration,  
Bangkapi District, Bangkok, 10240, Thailand

---

**Abstract: Problem statement:** For a multivariate normal population with size smaller than dimension,  $n < p$ , the likelihood ratio tests of the null hypothesis that the mean vector was zero with a one-sided alternative were no longer valid because they involved with sample covariance matrix which was singular. **Approach:** The test statistics for one-sided multivariate hypotheses with  $n < p$  were proposed. **Results:** The simulation study showed that the proposed tests provided reasonable type I error rate for one-sided covariance structures. They also give good powers. The application of these tests was given by testing of one-sided hypotheses on DNA micro array data. **Conclusion:** Under that there have no such other tests available at present for this kind of hypothesis testing with  $n < p$  yet, the proposed tests are good ones. However, the methodology is valid for any one-sided hypotheses application which involves high-dimensional data.

**Key words:** DNA micro arrays, multivariate normal, one-sided multivariate test, Follmann's test, power comparison

---

### INTRODUCTION

Suppose one uses a matched-pair design to compare the multivariate responses of two treatments. If the responses are  $p$  dimensional and  $\theta = (\theta_1, \theta_2, \dots, \theta_p)$  is the difference, treatment one minus treatment two, of the mean responses, then one may test the null hypothesis,  $H_0: \theta_1 = \theta_2 = \dots = \theta_p = 0$ , to determine if there is a difference in the two treatments. Furthermore, if one believes that for each coordinate, the mean responses for treatment one are at least as large as those for treatment two, then the alternative can be constrained by  $H_1: \theta_i \geq 0$  for  $i = 1, 2, \dots, p$ .

Based on a random sample with  $n > p$  from the normal distribution with mean  $\theta$  and covariance matrix  $V$ , Kudo (1963); Shorack (1967) and Perlman (1969) derived the likelihood ratio test of  $H_0$  versus  $H_1-H_0$  for the cases in which  $V$  is known, known up to a multiplicative constant and completely unknown, respectively. Because the likelihood ratio tests with restricted alternatives are complicated to use, Tang *et al.* (1989) proposed an approximate likelihood ratio test and Follmann (1996) proposed one-sided modifications of the usual  $\chi^2$  and Hotelling's  $T^2$  tests of  $H_0$  versus  $\sim H_0$  that are easier to implement. Using exact computations and Monte Carlo methods, Chongcharoen *et al.* (2002) compared the performance of Kudo's test, Follmann's test, a new test, which is a modification of Follmann's

test, the permutation test of Boyett and Shuster (1977) and the Tang-Gnecco-Geller test for a known covariance matrix and for a partially known covariance matrix, they compared the powers of these tests with Kudo's test replaced by Shorack's test. For a completely unknown covariance matrix, Chongcharoen (2009) studied the power of these one-sided tests for unknown covariance matrices with equal variances and unequal variances as well as tests obtained by combining the Boyett and Shuster (1977) technique to Follmann's test, the new test, Perlman's test and the Tang-Gnecco-Geller test.

In some situations, there are no longer data for  $n > p$ . That is, when the number  $n$  of available observations is smaller than the dimension  $P$  of the observed vectors. For example, the data come from DNA micro arrays where thousands of gene expression levels are measured in relatively few subjects. The one-sided multivariate tests as above are no longer valid for this kind of data because the  $p \times p$  sample covariance matrix  $S$  is singular with rank  $n < p$ ,  $S^{-1}$  does not exist. Since now there have no one-sided multivariate tests available for the data which has the number  $n$  of available observations is smaller than the dimension  $p$  yet, therefore the proposed tests were the one-sided multivariate tests for the data with  $n < p$ .

Throughout this study, suppose  $X_1, X_2, \dots, X_n$  is a random sample from a  $p$ -dimensional multivariate

normal distribution with unknown mean  $\theta = (\theta_1, \theta_2, \dots, \theta_p)'$  and unknown positive definite covariance matrix  $V$  with  $n \leq p$ . One may consider testing the null hypothesis  $H_0: \theta = 0$  versus  $H_1: H_0$  where  $H_1: \theta \in \Omega_p$  and  $\Omega_p = \{x | x_i \geq 0; i=1,2,\dots,p\}$  is the  $p$ -dimensional nonnegative orthant. The sample mean and covariance are Eq. 1:

$$\bar{X} = \sum_{i=1}^n \frac{X_i}{n} \quad \text{and} \quad S = \sum_{i=1}^n \frac{(X_i - \bar{X})(X_i - \bar{X})'}{n-1} \quad (1)$$

when  $n < p$ ,  $S$  is a singular matrix.

The hypotheses  $H_0$  and  $H_1$  also arise in the one-way analysis of variance when the means are known to satisfy an order restriction. For observations which come from  $k$  normal populations whose means are known to satisfy a simple ordering, i.e.,  $H_S: \mu_1 \leq \mu_2 \leq \dots \leq \mu_k$ , Bartholomew (1959; 1961) derived the likelihood ratio test of  $\mu_1 = \mu_2 = \dots = \mu_k$  with the alternative restricted by  $H_S$  for the cases of known variances and variances known up to a multiplicative constant. Suppose the observations of a random sample are  $Y_{ij}$  for  $j=1,2,\dots,n_i$  and  $i=1,2,\dots,k$  and the sample means are  $\bar{Y}_1, \bar{Y}_2, \dots, \bar{Y}_k$ . With known variances,  $\sigma_1^2, \sigma_2^2, \dots, \sigma_k^2$ , Kudo (1963) noted that for  $p=k-1$ ,  $X_\ell = \bar{Y}_{\ell+1} - \bar{Y}_\ell$  for  $\ell=1,2,\dots,p$ ,  $X = (X_1, X_2, \dots, X_p)'$  and  $\theta = E(X)$ , the hypotheses on  $\mu$  are equivalent to  $H_0$  and  $H_1$  above, Bartholomew's and Kudo's tests are equivalent. If the weights  $w_i = n_i / \sigma_i^2$  are equal, then the correlation matrix for  $X$  of simple order restriction is Eq. 2:

$$(R_S)_{\ell t} = I(\ell = t) - 0.5I(|t - \ell| = 1) \quad \text{for } 1 \leq \ell, t \leq p \quad (2)$$

where,  $I(A)$  is the indicator of  $A$ . Also, Bartholomew (1959; 1961) considered an arbitrary partial order restriction, which includes the simple tree order, i.e.,  $H_T: \mu_i \leq \mu_j$  for  $i=2,3,\dots,k$ . For this ordering, one takes differences,  $X_\ell = \bar{Y}_{\ell+1} - \bar{Y}_1$  for  $\ell=1,2,\dots,p$  and with  $p=k-1$  and  $w_i$  as above, the correlation matrix of  $X = (X_1, X_2, \dots, X_p)$  for simple order tree restriction is Eq. 3:

$$(R_T)_{\ell t} = I(\ell = t) + 0.5I(\ell \neq t) \quad \text{for } 1 \leq \ell, t \leq p \quad (3)$$

where,  $I(A)$  is the indicator of  $A$  as above. In this study, we mainly interested in one-sided multivariate tests which involved both  $R_S$  and  $R_T$ . So the powers of the proposed tests are compared for  $R_S$  and  $R_T$  including several other correlation matrices.

## MATERIALS AND METHODS

**The unrestricted alternative test for high dimensional multivariate tests:** The unrestricted alternative test for mean of random sample  $X_1, X_2, \dots, X_n$  with  $X_i \sim \text{iid}N(\theta, V)$  when  $n < p$ , that is, the tests with the hypothesis Eq. 4:

$$H_0: \theta_1 = \theta_2 = \dots = \theta_p = 0, H_1: \text{at least } \theta_i \neq 0 \quad (4)$$

are proposed by several researchers recently such as:

Dempster (1958; 1960) proposed a test for testing the mean difference of two independent samples and developed its approximate distribution. Srivastava (2007) and Srivastava and Du (2008) gave the one sample version of Dempster's test statistic which rejects  $H_0$  as (4) at a significant level  $\alpha$  if:

$$D = \frac{n\bar{X}'\bar{X}}{\text{tr}(S)} > F_{\alpha; [\hat{r}], [(n-1)\hat{r}]}$$

Where:

$\bar{X}$  = The sample mean vector  
 $S$  = The sample covariance defined as in (1)

$F_{\alpha; [\hat{r}], [(n-1)\hat{r}]}$  is the  $(1-\alpha)$  th quintile of the F-distribution with degrees of freedom  $[\hat{r}]$  and  $[(n-1)\hat{r}]$ , where  $[a]$  denotes the largest integer less than or equal to  $a$  and:

$$\hat{r} = p \frac{\hat{a}_1^2}{\hat{a}_2}, \quad \hat{a}_1 = \frac{\text{tr}(S)}{p} \quad \text{and} \quad \hat{a}_2 = \frac{(n-1)^2}{(n-2)(n+1)p} \left[ \text{tr}(S^2) - \frac{(\text{tr}(S))^2}{n-1} \right]$$

Under condition  $0 < \lim_{p \rightarrow \infty} a_i = \lim_{p \rightarrow \infty} \frac{\text{tr}(V^i)}{p} = a_{i0} < \infty;$

$i=1,2,3,4$ . This test is the uniformly most powerful test among all the test which are invariant under transformation  $X_i \rightarrow c\Gamma X_i$  where  $c \neq 0$  and  $\Gamma\Gamma' = I_p$ . They compared this version of Dempster's test with Bai-Saranadasa's test and their test which we will discuss after studying their test.

Bai and Saranadasa (1996) also proposed a test for testing the mean difference of two independent samples. They derived its asymptotic power of their test. Also they derived the asymptotic power of the classical Hotelling's  $T^2$  test and Dempster's non-exact test for a two-sample problem.

Table 1: Attained significance level of Dempster’s test (D), Bai and Saranadasa’s test (BS) and Srivastava and DU’s test(SD) under the null hypothesis for correlation matrix  $\mathfrak{R} = R_s$ ,  $\mathfrak{R} = R_r$ , with all off-diagonal elements equals to -0.5 called  $\mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2 = [\rho_{ij}]$ ,  $\rho_{ij} = -0.5$  for  $j=2i$  and  $\rho_{ij} = 0.7$  for elsewhere for  $1 \leq i, j \leq p$  at  $\alpha = 0.05$

p	n	$\mathfrak{R} = R_s$			$\mathfrak{R} = R_r$			$\mathfrak{R} = R_1$			$\mathfrak{R} = R_2$		
		D	BS	SD									
10	5	0.066	0.086	0.16	0.094	0.116	0.158	0.073	0.125	0.081	0.078	0.122	0.103
	10	0.049	0.058	0.094	0.069	0.084	0.090	0.053	0.089	0.050	0.054	0.084	0.058
20	10	0.051	0.055	0.096	0.075	0.088	0.086	0.055	0.096	0.038	0.052	0.088	0.042
	20	0.053	0.052	0.072	0.061	0.069	0.060	0.045	0.068	0.025	0.044	0.071	0.029
30	10	0.054	0.056	0.097	0.071	0.083	0.079	0.053	0.088	0.030	0.049	0.087	0.035
	15	0.053	0.052	0.080	0.062	0.072	0.062	0.047	0.074	0.021	0.044	0.073	0.023
	20	0.055	0.052	0.070	0.059	0.067	0.053	0.046	0.067	0.018	0.047	0.074	0.022
40	30	0.049	0.047	0.060	0.054	0.063	0.046	0.044	0.062	0.015	0.045	0.063	0.018
	10	0.054	0.057	0.096	0.076	0.084	0.080	0.053	0.094	0.027	0.053	0.089	0.030
	20	0.056	0.053	0.071	0.058	0.067	0.051	0.047	0.067	0.016	0.046	0.072	0.018
	30	0.051	0.047	0.062	0.054	0.062	0.043	0.047	0.064	0.012	0.048	0.064	0.015
50	40	0.050	0.045	0.054	0.055	0.060	0.042	0.048	0.060	0.011	0.047	0.060	0.014
	10	0.057	0.059	0.096	0.079	0.088	0.080	0.049	0.088	0.023	0.052	0.090	0.024
	20	0.056	0.052	0.068	0.065	0.073	0.054	0.049	0.069	0.013	0.050	0.074	0.016
	25	0.049	0.046	0.058	0.057	0.065	0.045	0.048	0.064	0.013	0.046	0.064	0.014
	30	0.049	0.046	0.056	0.058	0.065	0.044	0.048	0.065	0.012	0.049	0.065	0.014
60	40	0.052	0.048	0.057	0.054	0.060	0.039	0.049	0.062	0.010	0.046	0.060	0.012
	50	0.048	0.046	0.050	0.051	0.058	0.034	0.047	0.058	0.009	0.048	0.062	0.008
	10	0.054	0.053	0.095	0.075	0.083	0.075	0.053	0.087	0.022	0.052	0.085	0.025
	20	0.054	0.052	0.066	0.061	0.069	0.048	0.048	0.068	0.013	0.052	0.073	0.013
	30	0.050	0.046	0.057	0.054	0.062	0.038	0.047	0.063	0.009	0.046	0.060	0.011
	40	0.052	0.050	0.055	0.056	0.062	0.036	0.045	0.058	0.008	0.045	0.059	0.009
100	50	0.048	0.048	0.050	0.056	0.062	0.036	0.049	0.060	0.009	0.049	0.058	0.009
	60	0.054	0.052	0.056	0.055	0.062	0.035	0.050	0.059	0.008	0.048	0.059	0.009
	10	0.056	0.054	0.088	0.079	0.087	0.072	0.050	0.089	0.016	0.050	0.087	0.017
	20	0.053	0.050	0.056	0.064	0.072	0.046	0.051	0.070	0.009	0.051	0.071	0.010
	50	0.049	0.047	0.049	0.049	0.056	0.029	0.047	0.057	0.005	0.049	0.059	0.005
200	10	0.055	0.056	0.066	0.081	0.089	0.066	0.056	0.094	0.013	0.056	0.091	0.012
	20	0.051	0.049	0.048	0.062	0.070	0.036	0.051	0.069	0.006	0.052	0.071	0.007
	50	0.049	0.049	0.045	0.050	0.056	0.022	0.048	0.058	0.002	0.048	0.059	0.003
400	10	0.054	0.055	0.050	0.082	0.090	0.055	0.054	0.087	0.008	0.052	0.091	0.008
	20	0.053	0.053	0.036	0.063	0.070	0.027	0.054	0.070	0.004	0.051	0.070	0.004
	50	0.052	0.051	0.040	0.053	0.059	0.015	0.049	0.058	0.001	0.050	0.059	0.001

They compared their test with Dempster’s test and the classical Hotelling’s  $T^2$  test by a simulation study shown in Table 1 and 2 in their study which the results showed that both Dempster’s non-exact test and their test have higher power than the Hotelling’s  $T^2$  test when the data dimension is proportionally close to the sample size. They claimed that the power of Dempster’s test and their test are rather close but their test has always a higher power than those of Dempster’s tests. They also claimed that their test is still comparatively better than Dempster’s test because their test does not rely on the normality assumption. For normal cases and higher dimension the power of Dempster’s test and their test are almost the same. The reader should note that their simulation study showed only on the population

covariance form as  $\Sigma = I_p$  and  $\Sigma = (1-\rho)I_p + \rho J_p$  where  $J_p$  is a  $p \times p$  matrix with all entries one and  $\rho = 0.5$  for normal case and  $\rho = 0, 0.3, 0.6$  and  $0.9$  for non-normal case.

From the fact if the test statistic of the proposed test has corrected distribution, then the proportion of rejection the null hypothesis under the null hypothesis is true, or called attained significance level, from the simulation result must close to the probability rejecting the null hypothesis when the null hypothesis is true, or called significance level, here is  $\alpha$ , which in their study they set the target significance level as  $\alpha = 0.05$ . From Table 1 for non-normal case in their study, the attained significance level of their test is close to  $\alpha = 0.05$  with the maximum difference from the target 0.003.

Table 2: Attained significance level of DF and BSF under the null hypothesis when the covariance matrices are  $\mathfrak{R} = R_s, \mathfrak{R} = R_T, \mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2$ , respectively

p	n	$\mathfrak{R} = R_s$		$\mathfrak{R} = R_T$		$\mathfrak{R} = R_1$		$\mathfrak{R} = R_2$	
		DF	BSF	DF	BSF	DF	BSF	DF	BSF
10	5	0.056	0.060	0.067	0.069	0.058	0.072	0.063	0.070
	10	0.051	0.045	0.056	0.053	0.047	0.051	0.053	0.054
20	10	0.051	0.048	0.056	0.053	0.052	0.055	0.049	0.053
	20	0.053	0.048	0.049	0.043	0.050	0.046	0.043	0.041
30	10	0.054	0.051	0.054	0.049	0.051	0.053	0.049	0.052
	15	0.052	0.049	0.047	0.043	0.051	0.048	0.045	0.044
	20	0.056	0.051	0.047	0.042	0.051	0.044	0.047	0.043
40	30	0.047	0.041	0.046	0.040	0.047	0.038	0.046	0.040
	10	0.054	0.053	0.055	0.050	0.052	0.053	0.051	0.054
	20	0.055	0.050	0.047	0.041	0.050	0.044	0.046	0.041
50	30	0.051	0.044	0.047	0.040	0.049	0.041	0.048	0.040
	40	0.048	0.042	0.041	0.035	0.046	0.036	0.051	0.041
	10	0.055	0.052	0.057	0.053	0.051	0.053	0.051	0.053
	20	0.051	0.047	0.049	0.044	0.050	0.044	0.049	0.045
60	25	0.048	0.046	0.044	0.039	0.051	0.042	0.050	0.043
	30	0.049	0.045	0.045	0.039	0.050	0.042	0.048	0.039
	40	0.050	0.043	0.044	0.038	0.050	0.039	0.048	0.038
	50	0.049	0.047	0.040	0.034	0.047	0.038	0.049	0.041
	10	0.057	0.052	0.057	0.052	0.048	0.049	0.051	0.052
	20	0.050	0.047	0.050	0.043	0.055	0.047	0.049	0.043
100	30	0.046	0.046	0.045	0.039	0.050	0.040	0.047	0.038
	40	0.051	0.045	0.045	0.039	0.048	0.037	0.049	0.038
	50	0.051	0.048	0.044	0.038	0.052	0.042	0.047	0.035
	60	0.054	0.051	0.045	0.039	0.051	0.039	0.048	0.037
	10	0.054	0.051	0.058	0.054	0.053	0.054	0.051	0.052
	20	0.049	0.047	0.047	0.041	0.051	0.042	0.050	0.044
200	50	0.047	0.047	0.043	0.036	0.051	0.040	0.049	0.038
	10	0.052	0.052	0.059	0.054	0.054	0.055	0.053	0.054
	20	0.054	0.051	0.049	0.044	0.052	0.043	0.052	0.046
400	50	0.049	0.047	0.045	0.039	0.050	0.039	0.050	0.038
	10	0.052	0.054	0.058	0.053	0.055	0.056	0.051	0.051
	20	0.057	0.054	0.050	0.043	0.052	0.042	0.053	0.045
	50	0.050	0.050	0.044	0.037	0.052	0.040	0.049	0.036

For normal cases as in Table 2 in their study, the attained significance level of their test is not close to 0.05 with the minimum difference from the target 0.012 meanwhile the attained significance level of Dempster’s test is also close to the target for non-normal case but it is equal to 0.05 for normal case with only  $\Sigma = I_p$ . It is true that all the attained significance level results showed in their study are within the range 0.0316 as mentioned in their study but we may be looking for a better test which gives the attained significance level closer to the target significance level. From this simulation results, it may possible that both their test and Dempster’s test can be used only some covariance matrix structure cases and they may need to be studied further. Srivastava (2007) and Srivastava and Du (2008) also gave the one sample version of Bai and Saranadasa’s test statistic which reject  $H_0$  as (4) at a significant level  $\alpha$  if:

$$BS > z_{\frac{\alpha}{2}} \text{ or } BS < -z_{\frac{\alpha}{2}}$$

Where:

$$BS = \frac{n\bar{X}'\bar{X} - \text{tr}(S)}{\left[ \frac{n}{n-1} \right]^2 \left[ 2p\hat{a}_2 \right]^2} = \frac{n\bar{X}'\bar{X} - \text{tr}(S)}{\left[ \frac{2n(n-1)}{(n-2)(n+1)} (\text{tr}(S^2)) \right]^2 - \frac{(\text{tr}(S))^2}{n-1}}$$

and  $z_{\frac{\alpha}{2}}$  is the  $(1 - \frac{\alpha}{2})$  Th quintile of the standard normal distribution and under conditions

$$0 < \lim_{p \rightarrow \infty} a_i = \lim_{p \rightarrow \infty} \frac{(\text{tr}(V^i))}{p} = a_{i0} < \infty, i = 1, 2, 3, 4 \text{ and for}$$

$\lambda_i = O(p^\gamma), 0 \leq \gamma \leq \frac{1}{2}$ , where  $\lambda_i$  are the eigenvalues of V.

Under the null hypothesis:

$$\lim_{(n,p) \rightarrow \infty} P_0(BS \leq z) = \Phi(z)$$

This test is also invariant under transformation  $X_i \rightarrow c\Gamma X_i$  where  $c \neq 0, \Gamma\Gamma' = I_p$  as Dempster's test does.

Srivastava (2007) and Srivastava and Du (2008) proposed a test for one sample which is based on the test statistic:

$$SD = \frac{n\bar{X}'D_s^{-1}\bar{X} - \frac{(n-1)p}{(n-3)}}{\sqrt{2(\text{tr}(R^2) - \frac{p^2}{n-1})(1 + \frac{\text{tr}(R^2)}{p^{\frac{2}{3}}})}}$$

where,  $R = D_s^{-\frac{1}{2}}SD_s^{-\frac{1}{2}}$  is the sample correlation matrix and  $D = \text{diag}(s_{11}, \dots, s_{pp})$  is the diagonal matrix with the diagonal elements of  $S$  defined in (1). Under conditions stated in their study, when  $\theta = 0$ ,  $SD$  is asymptotically distributed as  $N(0, 1)$ . Then this test will reject  $H_0$  as (4) at a significant level  $\alpha$  if:

$$SD > z_{\frac{\alpha}{2}} \text{ or } SD < -z_{\frac{\alpha}{2}}$$

where,  $z_{\frac{\alpha}{2}}$  is the  $(1 - \frac{\alpha}{2})$ th quantile of the standard normal distribution. They also showed that this test is an invariant test under the group of scalar transformations  $X_i \rightarrow TX_i$ , where  $T = \text{diag}(t_1, t_2, \dots, t_p)$  and  $t_1, t_2, \dots, t_p$  are nonzero constants. They claimed by simulation that for all the components of the random vector are independent, that is, the covariance matrix is a diagonal matrix, their test has the attained significance level given in Table 1 in their study reasonably well in all cases. But we can see in Table 1 in their study that all attained significance level values vary from 0.035-0.065. There is a number of attaining significance level values differ from 0.05. They also claimed that their test has substantial better power than Dempsters's test and Bai and Saranadasa's test which showed only on diagonal covariance structures. They did not show the attained significance level of their test for other covariance structures. One may have questions about the power of these tests for other forms of covariance structures.

So, we will investigate these tests, Dumpster's test, Bai and Saranadasa's test and Srivastava and Du's test, for one sample in the other forms of covariance structures mainly on the covariance from the restricted alternative hypotheses,  $H_0 : \theta = 0$  versus  $H_1 - H_0$  where  $H_1 : \theta \in \Omega_p$  and  $\Omega_p = \{x | x_i \geq 0; i = 1, 2, \dots, p\}$ , that is,  $R_S$  and  $R_T$  as well as the correlation matrix with all off-diagonal elements equals to -0.5 called  $\Re = R_1$ , and the correlation matrix  $\Re = R_2 = [\rho_{ij}]$ ,  $\rho_{ij} = -0.5$  for  $j=2i$  and

$\rho_{ij} = 0.7$  for elsewhere. Their estimated significant levels are computed by using the Monte Carlo method for:

$p = 10, n = 5, 10; p = 20, n = 10, 20; p = 30, n = 10, 15, 20, 30; p = 40, n = 10, 20, 30, 40; p = 50, n = 10, 20, 30, 40, 50; p = 60, n = 10, 20, 30, 40, 50, 60; p = 100, n = 10, 20, 50; p = 200, n = 10, 20, 50; p = 400, n = 10, 20, 50$ . Each case is repeated 10,000 times and the proportion of rejections record for each test. All of these tests are conducted using the level of significance  $\alpha = 0.05$ . It was found in Table 1 that with their critical values, the estimated significance level of these tests, except  $D$  test statistic with  $R_S$ , under all covariance matrices considered, including  $R_S$  and  $R_T$ , are not consistent in  $\alpha = 0.05$ . That is, the estimated significance level of some tests is too large with some covariance matrices meanwhile the other tests give the estimated significance level too small with the other covariance structure considered. For instance, the estimated significance level for  $D$ 's test approximately close to  $\alpha = 0.05$  very well for each covariance structure and each  $p, n$  considered, the  $BS$ 's test gave the estimated significance level reasonably well only on the simple order for each  $p, n$  and gave the poor estimated significance level for other forms of covariance considered and the  $SD$ 's test gave an estimated significance level which do not consist to  $\alpha = 0.05$ . Since  $BS$ 's test and  $SD$ 's test gave the poor estimated significance level for non-diagonal covariance matrix, it is shown that these two tests may not suitable for high-dimensional data with non-diagonal covariance matrix. But  $BS$ 's test is at least well on simple order covariance matrix and also Bai and Saranadasa (1996) showed that their test,  $BS$ 's test, has asymptotic powers the same as those of  $D$ 's test, thus we will investigate the  $D$ 's test and  $BS$ 's test further for one-side alternatives.

**The restricted alternative proposed tests for high dimensional multivariate tests:** For the tests with restricted alternatives, that is, to test the null hypothesis  $H_0 : \theta = 0$  versus  $H_1 - H_0$  where  $H_1 : \theta \in \Omega_p$  and  $\Omega_p = \{x | x_i \geq 0; i = 1, 2, \dots, p\}$ , one may apply (Follmann, 1996) test to both  $D$ 's test and  $BS$ 's test. When applied to  $D$ 's test which is denoted  $DF$ , it rejects  $H_0$  at level  $\alpha$  if:

$$D > F_{2\alpha; [r], [(n-1)r]} \text{ and } \sum_{j=1}^p \bar{X}_j > 0$$

where,  $F_{2\alpha; [r], [(n-1)r]}$  is the  $(1 - 2\alpha)$ th quantile of the central  $F$ -distribution with  $[r]$  and  $[(n-1)r]$  degrees of

freedom. By Theorem 2.1 of Follmann (1996), one has  $\theta = 0$  and the significance level is approximated by:

$$\begin{aligned} & \Pr(D > F_{1-2\alpha, [r], [(n-1)r]} \cap \bar{1}'\bar{X} > 0) \\ &= \Pr(D > F_{1-2\alpha, [r], [(n-1)r]}) \Pr(\bar{1}'\bar{X} > 0) \\ &= (2\alpha) \times \frac{1}{2} \\ &= \alpha \end{aligned}$$

Also, when applied Follmann's idea to BS's test, called BSF, one may reject  $H_0$  at level  $\alpha$  if:

$$BS > z_\alpha \text{ or } BS < -z_\alpha$$

And:

$$\sum_{j=1}^p \bar{X}_j > 0$$

where,  $Z_\alpha$  is the  $(1-\alpha)^{\text{th}}$  quantile of the standard normal distribution. It also noted that, after Theorem 2.1 of Follmann (1996), the significance level of this test is  $\alpha$ .

In Table 2., for every  $p$  and  $n > 5$  considered, DF gives the estimated significance level range from 0.047-0.057 for RS, range from 0.044-0.058 for the RT, range from 0.047-0.055 for the correlation matrix  $\mathfrak{R} = R_1$  and range from 0.043-0.053 for  $\mathfrak{R} = R_2$  and BSF gives the estimated significance level range from 0.041-0.054 for RS, range from 0.036-0.054 for the  $R_T$ , range from 0.036-0.056 for the correlation matrix  $\mathfrak{R} = R_1$  and range from 0.036-0.054 for  $\mathfrak{R} = R_2$ . It is shown that the estimated significance levels of both tests approximate reasonably well in all cases considered.

Table 3: Comparison of power of D and DF under the alternative hypothesis when the covariance matrices are  $\mathfrak{R} = R_S, \mathfrak{R} = R_T, \mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2$ , respectively

p	n	$\mathfrak{R} = R_S$		$\mathfrak{R} = R_T$		$\mathfrak{R} = R_1$		$\mathfrak{R} = R_2$	
		D	DF	D	DF	D	DF	D	DF
10	5	0.552	0.721	0.389	0.475	0.231	0.328	0.166	0.227
	10	0.984	0.998	0.635	0.746	0.389	0.593	0.239	0.357
20	10	0.996	0.999	0.563	0.675	0.271	0.423	0.149	0.263
	20	1.000	1.000	0.982	0.994	0.824	0.942	0.428	0.603
30	10	1.000	1.000	0.769	0.852	0.440	0.625	0.216	0.337
	15	1.000	1.000	0.869	0.932	0.586	0.758	0.283	0.419
	20	1.000	1.000	0.968	0.991	0.781	0.939	0.347	0.509
40	30	1.000	1.000	1.000	1.000	0.982	0.998	0.623	0.784
	10	1.000	1.000	0.730	0.826	0.387	0.558	0.190	0.301
	20	1.000	1.000	0.952	0.983	0.694	0.859	0.322	0.471
	30	1.000	1.000	0.996	0.999	0.929	0.988	0.517	0.676
50	40	1.000	1.000	1.000	1.000	0.969	0.997	0.630	0.776
	10	1.000	1.000	0.740	0.833	0.390	0.544	0.185	0.303
	20	1.000	1.000	0.987	0.996	0.797	0.921	0.407	0.567
	25	1.000	1.000	0.999	1.000	0.941	0.988	0.548	0.702
	30	1.000	1.000	0.998	0.999	0.913	0.975	0.526	0.679
60	40	1.000	1.000	1.000	1.000	0.996	1.000	0.745	0.868
	50	1.000	1.000	1.000	1.000	0.997	0.999	0.749	0.886
	10	1.000	1.000	0.744	0.842	0.380	0.556	0.178	0.286
	20	1.000	1.000	0.987	0.996	0.780	0.904	0.390	0.549
	30	1.000	1.000	0.994	0.999	0.876	0.968	0.446	0.612
	40	1.000	1.000	1.000	1.000	0.994	0.999	0.746	0.864
100	50	1.000	1.000	1.000	1.000	0.998	1.000	0.784	0.894
	60	1.000	1.000	1.000	1.000	1.000	1.000	0.890	0.969
	10	1.000	1.000	0.785	0.871	0.428	0.589	0.216	0.328
	20	1.000	1.000	0.975	0.992	0.735	0.872	0.354	0.501
	50	1.000	1.000	1.000	1.000	0.999	1.000	0.759	0.886
200	10	1.000	1.000	0.753	0.844	0.391	0.545	0.198	0.303
	20	1.000	1.000	0.976	0.993	0.731	0.863	0.357	0.497
	50	1.000	1.000	1.000	1.000	0.997	1.000	0.745	0.872
400	10	1.000	1.000	0.768	0.849	0.405	0.56	0.195	0.299
	20	1.000	1.000	0.971	0.992	0.699	0.845	0.333	0.475
	50	1.000	1.000	1.000	1.000	0.999	1.000	0.787	0.904

Table 4: Comparison of power of BS and BSF under the alternative hypothesis when the covariance matrices are  $\mathfrak{R} = R_S, \mathfrak{R} = R_T, \mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2$ , respectively

p	n	$\mathfrak{R} = R_S$		$\mathfrak{R} = R_T$		$\mathfrak{R} = R_1$		$\mathfrak{R} = R_2$	
		BS	BSF	BS	BSF	BS	BSF	BS	BSF
10	5	0.643	0.718	0.456	0.488	0.368	0.383	0.247	0.252
	10	0.991	0.998	0.698	0.737	0.566	0.621	0.332	0.367
20	10	0.996	0.998	0.619	0.663	0.401	0.440	0.229	0.249
	20	1.000	1.000	0.989	0.993	0.893	0.929	0.534	0.591
30	10	1.000	1.000	0.813	0.847	0.587	0.638	0.315	0.349
	15	1.000	1.000	0.897	0.924	0.692	0.743	0.367	0.410
	20	1.000	1.000	0.981	0.988	0.877	0.918	0.436	0.485
40	30	1.000	1.000	1.000	1.000	0.992	0.996	0.701	0.752
	10	1.000	1.000	0.780	0.816	0.524	0.569	0.282	0.312
	20	1.000	1.000	0.968	0.979	0.780	0.829	0.401	0.445
	30	1.000	1.000	0.998	0.999	0.958	0.975	0.583	0.635
50	40	1.000	1.000	1.000	1.000	0.983	0.993	0.683	0.732
	10	1.000	1.000	0.785	0.823	0.508	0.550	0.280	0.311
	20	1.000	1.000	0.993	0.996	0.861	0.897	0.486	0.535
	25	1.000	1.000	0.999	1.000	0.968	0.981	0.617	0.668
	30	1.000	1.000	0.999	0.999	0.940	0.960	0.588	0.639
60	40	1.000	1.000	1.000	1.000	0.998	0.999	0.789	0.832
	50	1.000	1.000	1.000	1.000	0.999	0.999	0.795	0.839
	10	1.000	1.000	0.794	0.832	0.518	0.565	0.267	0.295
	20	1.000	1.000	0.993	0.995	0.844	0.882	0.462	0.513
	30	1.000	1.000	0.998	0.998	0.915	0.946	0.509	0.564
	40	1.000	1.000	1.000	1.000	0.997	0.999	0.788	0.828
100	50	1.000	1.000	1.000	1.000	0.999	1.000	0.820	0.859
	60	1.000	1.000	1.000	1.000	1.000	1.000	0.915	0.944
	10	1.000	1.000	0.830	0.862	0.548	0.594	0.301	0.333
	20	1.000	1.000	0.984	0.990	0.801	0.845	0.420	0.465
	50	1.000	1.000	1.000	1.000	1.000	1.000	0.796	0.840
200	10	1.000	1.000	0.797	0.833	0.505	0.549	0.276	0.307
	20	1.000	1.000	0.985	0.991	0.795	0.837	0.417	0.464
	50	1.000	1.000	1.000	1.000	0.998	0.999	0.779	0.826
400	10	1.000	1.000	0.806	0.841	0.519	0.561	0.270	0.301
	20	1.000	1.000	0.983	0.989	0.769	0.814	0.391	0.438
	50	1.000	1.000	1.000	1.000	1.000	1.000	0.815	0.856

**RESULTS**

To compare these two tests, the performances of them are studied by Monte Carlo techniques for multivariate normal distributions with the correlation matrices  $R_S$  and  $R_T$ , that is for the simple order and the simple tree order correlations with equal weights as well as some other forms of correlation structures such as  $\mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2$ . Recall,  $R_S$  and  $R_T$  are given in (2) and (3), respectively. The mean vector for the alternative hypothesis is chosen in the non-negative orthant as  $\theta = (v_1, v_2, \dots, v_p)'$ ;  $v_{2k-1} = 0$  and  $v_{2k} \sim \text{iid Unif}(0,1), k = 1, 2, \dots, p/2$  so that the tests will be rejected. As before, 10,000 iterations are used. In each iteration,  $n$  multivariate normal  $X$ 's with the chosen mean vector and covariance of the form  $\mathfrak{R}$  are generated and the proportion of rejections for these tests was recorded. All of these tests are conducted using the level of

significance  $\alpha = 0.05$ . Monte-Carlo estimated power of these two tests is given in Table 3-5.

Table 3 gives the powers of the D's test and DF's test in all cases considered. It can be seen that DF's test gives at least better powers than D's test does. Also DF's test gives highest power when  $p, n$  large and  $p \gg n$  for  $R_S$ . Similar to comparison powers between BS's test and BSF's test, shown in Table 4. BSF's test showed substantially higher power than BS's test does in all cases considered and also gives highest power for  $R_S$  in each  $p$  and  $n > 5$ . To compare the BSF's test to DF's test in all cases considered, Table 5 gives their powers. For the correlation matrices  $R_S$  and  $R_T$ , both BSF's test and DF's test almost give the same powers and for correlation matrices as  $\mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2$ , for each  $p$  DF's test gives higher power than the BSF's test when  $n > 10$ . Therefore, we can conclude that overall both tests, BSF's test and DF's test, gave almost the same powers in every  $p$  and  $n$  and every covariance matrices structure considered.

Table 5: Empirical powers of DF and BSF under the alternative hypothesis when the covariance matrices are  $\mathfrak{R} = R_s, \mathfrak{R} = R_T, \mathfrak{R} = R_1$  and  $\mathfrak{R} = R_2$ , respectively.

p	n	$\mathfrak{R} = R_s$		$\mathfrak{R} = R_T$		$\mathfrak{R} = R_1$		$\mathfrak{R} = R_2$	
		DF	BSF	DF	BSF	DF	BSF	DF	BSF
10	5	0.721	0.718	0.475	0.488	0.328	0.383	0.227	0.252
	10	0.998	0.998	0.746	0.737	0.593	0.621	0.357	0.367
20	10	0.999	0.998	0.675	0.663	0.423	0.44	0.236	0.249
	20	1.000	1.000	0.994	0.993	0.942	0.929	0.603	0.591
30	10	1.000	1.000	0.852	0.847	0.625	0.638	0.337	0.349
	15	1.000	1.000	0.932	0.924	0.758	0.743	0.419	0.410
	20	1.000	1.000	0.991	0.988	0.939	0.918	0.509	0.485
40	30	1.000	1.000	1.000	1.000	0.998	0.996	0.784	0.752
	10	1.000	1.000	0.826	0.816	0.558	0.569	0.301	0.312
	20	1.000	1.000	0.983	0.979	0.859	0.829	0.471	0.445
50	30	1.000	1.000	0.999	0.999	0.988	0.975	0.676	0.635
	40	1.000	1.000	1.000	1.000	0.997	0.993	0.776	0.732
	10	1.000	1.000	0.833	0.823	0.544	0.550	0.303	0.311
	20	1.000	1.000	0.996	0.996	0.921	0.897	0.567	0.535
60	25	1.000	1.000	1.000	1.000	0.988	0.981	0.702	0.668
	30	1.000	1.000	0.999	0.999	0.975	0.960	0.679	0.639
	40	1.000	1.000	1.000	1.000	1.000	0.999	0.868	0.832
	50	1.000	1.000	1.000	1.000	0.999	0.999	0.886	0.839
	10	1.000	1.000	0.842	0.832	0.556	0.565	0.286	0.295
	20	1.000	1.000	0.996	0.995	0.904	0.884	0.549	0.513
100	30	1.000	1.000	0.999	0.998	0.968	0.946	0.612	0.564
	40	1.000	1.000	1.000	1.000	0.999	0.999	0.864	0.828
	50	1.000	1.000	1.000	1.000	1.000	1.000	0.894	0.859
	60	1.000	1.000	1.000	1.000	1.000	1.000	0.969	0.944
200	10	1.000	1.000	0.871	0.862	0.589	0.594	0.328	0.333
	20	1.000	1.000	0.992	0.990	0.872	0.845	0.501	0.465
	50	1.000	1.000	1.000	1.000	1.000	1.000	0.886	0.840
400	10	1.000	1.000	0.844	0.833	0.545	0.549	0.303	0.307
	20	1.000	1.000	0.993	0.991	0.863	0.837	0.497	0.464
	50	1.000	1.000	1.000	1.000	1.000	0.999	0.872	0.826
400	10	1.000	1.000	0.849	0.841	0.560	0.561	0.299	0.301
	20	1.000	1.000	0.992	0.989	0.845	0.814	0.475	0.438
	50	1.000	1.000	1.000	1.000	1.000	1.000	0.904	0.856

Table 6: Observed p-values for testing the changes in gene expression after treatment for leukemia data

	DF	BSF
<b>Leukemia data</b>		
Statistic	115.733	10.7072
Average sum	438022	438022
p-values	0.0129	0.0000

Therefore, for protection some gains of using these two tests, we recommend these tests for high dimensional data when  $p \geq 20$  and  $n > 10$  for one-side alternatives.

**An example:** The proposed tests are applied to an example of DNA micro array data which the data are 8280 (p) gene expression information on 110 childhoods suffering from acute lymphoblastic leukemia. To see the changes in gene expression after treatment, the data were cleaned and then obtained the difference of gene expression from before and after treatment of 50 children in 254 (p) gene expressions (<http://www.aialab.si/supp/bi-cancer/projections/info>).

The results of using these two tests are shown in Table 6. The p-values of DF's test and BSF's test equal to 0.0129, 0.0000 respectively. Thus, all two tests lead to the rejection of the hypothesis that the gene expressions after treatment have the same level as before treatment.

### DISCUSSION

At present, there have no such other tests available for this kind of hypothesis testing on high-dimensional data yet, the proposed test should be the best one available though it works well on some conditions or under the circumstances considered in this study. Hopefully, there will be some other researchers interested in it.

### CONCLUSION

Since for the data with the number n of available observations is smaller than the dimension p ( $n \leq p$ ), the

proposed one-sided multivariate tests, DF's test and BSF's test, have power larger than the tests with unrestricted multivariate alternative tests. Thus, for comparing the two treatments of data with the dimension  $p$  larger than the number  $n$  of available observations that one believes that for each coordinate the mean responses for treatment one are at least as large as those for treatment to which at present there have no such other tests available for this kind of hypothesis testing yet, we recommended the proposed tests, DF's test and BSF's test for  $p \geq 20$  and  $n > 10$  under the circumstances considered in this study.

#### ACKNOWLEDGEMENT

The researchers would like to express sincere thanks to Research center, The National institute of Development Administration, Bangkok, Thailand for financial support.

#### REFERENCES

- Bartholomew, D.J., 1959. A test of homogeneity for ordered alternatives. *Biometrika*, 46: 36-48. DOI: 10.1093/Biomet/46.1-2.36
- Bartholomew, D.J., 1961. A test of homogeneity of means under restricted alternatives. *J. Royal Stat. Soc. B. (Methodol.)*, 23: 239-281.
- Bai, Z. and H. Saranadasa, 1996. Effect of high dimension: By an example of a two sample problem. *Stat. Sinica.*, 6, 311-329.
- Boyett, J.M. and J.J. Shuster, 1977. Nonparametric one-sided tests in multivariate analysis with medical applications. *J. Am. Stat. Assoc.*, 72: 665-668.
- Chongcharoen, S., B. Singh and F.T. Wright, 2002. Powers of some one-sided multivariate tests with the population covariance matrix known up to a multiplicative constant. *J. Stat. Plann. Inference*, 107: 103-121. DOI: 10.1016/S0378-3758(02)00246-X
- Chongcharoen, S., 2009. Powers of some one-sided multivariate tests with unknown population covariance matrix. *Songklanakarin J. Sci. Technol.*, 31: 351-359.
- Dempster, A.P., 1958. A high dimensional two sample significance test. *Ann. Math. Stat.*, 29: 995-1010. DOI: 10.1214/aoms/1177706437
- Dempster, A.P., 1960. A significance test for the separation of two highly multivariate small samples. *Biometrics*, 16: 41-50.
- Follmann, D., 1996. A simple multivariate test for one-sided alternatives. *J. Am. Stat. Assoc.*, 91: 854-861. DOI: 10.2307/2291680
- Kudo, A., 1963. A multivariate analogue of the one-sided test. *Biometrika*, 50: 403-418. DOI: 10.1093/biomet/50.3-4.403
- Perlman, M.D., 1969. One-sided testing problems in multivariate analysis. *Ann. Math. Stat.*, 40: 549-567.
- Shorack, G.R., 1967. Testing against ordered alternatives in model I analysis of variance; normal theory and nonparametric. *Ann. Math. Stat.*, 38: 1740-1753. DOI: 10.1214/aoms/1177698608
- Srivastava, M.S., 2007. Multivariate theory for analyzing high dimensional data. *J. Japan Stat. Assoc.*, 37: 53-86.
- Srivastava, M.S. and M. Du, 2008. A test for the mean vector with fewer observations than the dimension. *J. Multivariate Anal.*, 99: 386-402. DOI: 10.1016/j.jmva.2006.11.002
- Tang, D.I., C. Gnecco and N.L. Geller, 1989. An approximate likelihood ratio test for a normal mean vector with nonnegative components with application to clinical trials. *Biometrika*, 76: 577-583. DOI: 10.1093/biomet/76.3.577